



Figure A.3.84: Percent of Families with Newborns Where the Mother Experienced Depression by District



Source: Manitoba Centre for Health Policy, 2008

Figure A.3.85: Percent of Families with Newborns Where the Mother Experienced Depression by Winnipeg Neighbourhood Clusters



Crude percent Source: Manitoba Centre for Health Policy, 2008

Table A.3.23: Crude Percent of Maternal Depression

Region	Crude Percent of Maternal Alcohol/Drug/Substance Use During Pregnancy	
	Number Observed per Year	CRUDE rate percent
	2000-2002	
South Eastman	20.7	3.5%
Central	36.3	3.5%
Assiniboine	24.7	4.9%
Brandon	19.0	3.8%
Winnipeg	291.0	4.8%
Interlake	32.3	6.5%
North Eastman	8.7	3.3%
Parkland	11.0	3.5%
Churchill	0.0	0.0%
Nor-Man	10.3	5.4%
Burntwood	11.7	4.0%
South	81.7	3.8%
Mid	52.0	4.8%
North	22.0	4.5%
Manitoba	466.7	4.5%

blank cells = suppressed


Source: Manitoba Centre for Health Policy, 2008

Region	Crude Percent of Maternal Alcohol/Drug/Substance Use During Pregnancy	
	Number Observed per Year	CRUDE rate percent
	2000-2002	
Fort Garry	20.3	3.8%
Assiniboine South	11.7	4.6%
St. Boniface	23.7	5.1%
St. Vital	31.7	5.6%
Transcona	11.7	3.5%
River Heights	22.3	4.7%
River East	36.7	4.3%
Seven Oaks	23.7	5.0%
St. James - Assiniboia	20.3	4.6%
Inkster	11.3	3.7%
Downtown	43.7	5.0%
Point Douglas	34.0	7.0%
Winnipeg	291.0	4.8%

blank cells = suppressed

Source: Manitoba Centre for Health Policy, 2008

**Figure A.3.86: Percent of Families with Newborns
Where the Mother Experienced Depression
by Aggregate RHA Area**



Source: Manitoba Centre for Health Policy, 2008